

Set Theoretical Forcing in Quantum Mechanics and AdS/CFT correspondence.

Jerzy Król^{‡§}

Abstract. We show unexpected connection of Set Theoretical Forcing with Quantum Mechanical lattice of projections over some separable Hilbert space. The basic ingredient of the construction is the rule of indistinguishability of Standard and some Nonstandard models of Peano Arithmetic. The ingeneric reals introduced by M. Ozawa will correspond to simultaneous measurement of incompatible observables. We also discuss some results concerning model theoretical analysis of Small Exotic Smooth Structures on topological 4-space $\mathbb{R}^4$. Forcing appears rather naturally in this context and the rule of indistinguishability is crucial again. As an unexpected application we are able to approach Maldacena Conjecture on *AdS/CFT* correspondence in the case of $AdS_5 \times S^5$ and Super YM Conformal Field Theory in 4 dimensions. We conjecture that there is possibility of breaking Supersymetry via sources of gravity generated in 4 dimensions by exotic smooth structures on $\mathbb{R}^4$ emerging in this context.

Keywords: Boolean Models of ZFC, Quantum Mechanics, Forcing, Exotic R^4 .

1. Remarks on Model Theory and QM.

Since the very beginning of Model Theory (MT), it was realized what is the position of formal language in the correct description of any Mathematical structure. There is a big difference when we talk about same things using different formal languages. Many "paradoxical" situations emerged in the development of the so called first order logic (let me mention only Löwenheim-Skolem Theorem or Gödel Theorems), which nonetheless serves as the logic depending on very minimal theoretical support from Set Theory and the like. It has resulted in emergence of various spectra of non isomorphic models, which first order theories must have (Keisler et al., 1990). One can say, that these are merely formal peculiarities, which are not acting upon our every day practice (espacially as a physicists). Surprisingly enough, the theories such as: Peano Arithmetic (PA), or axiomatized theory of natural numbers and Zermello-Fraenkel Set Theory (with the Axiom of Choice [AC]) ZF(C), or axiomatized theory of sets—both are first order theories. This means that all paradoxical situations concern also them. Because of the fact

[‡] Institute of Physics, University of Silesia, Uniwersytecka 4, 40-007 Katowice, Poland

[§] (iriking@poczta.fm)



that very effective formalism of Quantum Mechanics (QM) was developed very early, people were not considering tools of MT in the context of QM (also, they simply were not able to do so, because MT was being developed in the time Quantum Theory was as a young branch of Mathematics). In the paradigm of QM people found it possible to talk about Many Worlds (as in the Everett interpretation of QM), but no one has tried to incorporate Model Theoretical tools in the shape of pluralities of models of basic enough theory as ZFC is. In what follows we are giving some insights coming from MT, which might be helpful in analysing certain interpretational and formal aspects of Quantum Theory.

Firstly, in the face of the Löwenheim-Skolem Theorem (Keisler et al., 1990) it follows that ZFC (if it is consistent, Jech (1978)) has a countable model. In particular, the set of real numbers in this model is countable from the "outside" (Jech, 1978). Might it be, that discreteness of spectra of some physical Quantum Mechanical observables is connected to the discreteness of real numbers in some models of ZFC?

Secondly, might it be that pluralities and non isomorphism of models of ZF(C) (or some other formal theories) can be connected to pluralities of Quantum Mechanical worlds in Everettian style?

Thirdly, as follows from Kochen - Specker theorem (Kochen et al., 1967) and EPR kind of experiments, some physical observables cannot have definite values before measurements. Might this phenomenon be connected to nonexistence of some real numbers in some models of ZF(C)? Might it be that the real numbers in question, which are not present in one model, will appear in the other as a result of Set Theoretical forcing? That would mean that the reals obtained as a result of measuring of some observables would emerge as a result of being forced to belong to some enlarged universum of sets. In this situation it would mean that there are inherent reasons for considering a non- constant universum of sets in the context of QM.

Whereas we avoid deciding about the import of the first two situations, we are to explore the third one. In general, detailed proofs are not included here. Informal, intuitive explanations are often used in place of them. The proofs will be given elsewhere (Król, 2002).

We assume some elementary knowledge of Set Theoretical forcing; the general reference is Jech (1978) which also serves as a source for all set theoretical questions considered here. Some knowledge of Boolean Valued models of ZFC is also desirable; the general reference is Bell (1985). In the context of QM, Boolean Valued models are discussed in Takeuti (1978; 1979; 1983). General references concerning Topos Theory are Johnstone (1977) and MacLane et al. (1992).

There were some attempts recently, to interpret QM in terms of Topos Theory: Isham et al. (1999), Fearins (2002) (by the use of so called Smooth Toposes from Synthetic Differential Geometry (SDG) (Moerdijk et al., 1991)) or Raptis (2001) in the context of Quantum Gravity. Also, there are some attempts to approach QM through Model Theoretic tools (Benioff (2001), Król (2001), Takeuti (1978; 1979)). Much effort has been spent on working out connections of QM (and other branches of physics) with Nonstandard Analysis (Albeverio et al., 1986) invented by Robinson (1966) as a purely Model Theoretical task (Robinson himself was one of the creators and main contributors to Model Theory). Close connections of QM and formal logic were noticed from the very beginning of the subject (Neumann et al., 1936). The so called Quantum Logic (QL) emerged as a special one and different from Classical Logic and Intuitionistic Logic (Dalla Chiara et al., 2001).

Synthetic Differential Geometry (SDG) was worked out by Kock (1981) and then developed by Moerdijk et al. (1991). The aim of this approach was to build common framework for Toposes and Nonstandard Analysis in the Robinsonian style, as well as for differential geometry, which deals with infinitesimals called noninvertible or idempotent. Needless to say, such unification is of great interest to physicists, too. Because of that, there were some trials to use these techniques in the context of Quantum and Classical Gravity (Guts et al., 1999, 2001) or in QM (Fearins, 2002). Very nice features of Toposes, which are inducing Intuitionistic Logic, and are natural models for arbitrarily high order logic, cause people naturally start to think about using the strangeness of the world generated by Toposes (the Middle Third Law can be violated in Toposes, Standard Natural Numbers may be not a decidable object, there may not exist an object of naturals at all, Set Theory is generated in the inside of the Topos; Toposes might serve as Universes of Discourse for Mathematics, and so on (MacLane et al., 1992)) to the strangeness (at the interpretational level) of the Quantum World. However, one should be aware that:

Warning: *No Topos is able to interpret QM in the sense that the Logic generated by the Topos is the one generated by Quantum Mechanical phenomena.*

Proof: Logical structure of any Topos is based on Heyting Algebras (HA). Any HA is distributive one. Quantum logic is based on the Lattice of projections (Bell, 1985) in some separable Hilbert space. Such a Lattice is in general non distributive (Sikorski et al., 1963, Takeuti, 1978) (the lattice of projections is distributive only in the case of one dimensional Hilbert space). $\square$

2. Forcing, Set Theory and QM.

Forcing was discovered originally as a tool for proving some independence results in Set Theory from the axioms of $\text{ZF}(\text{C})$ (Cohen, 1963) and then, developed as very basic phenomenon in many different mathematical situations. Solovay and Scott (1969) and, independently, Vopenka (1965) have given formulation of forcing in the language of so called Boolean valued universes of $\text{ZF}(\text{C})$. Forcing as it stands has been shown to have specific significance in very general situations in the Model Theory (Robinson et al., 1971). Semantics connected to Toposes (at least to Grothendieck Toposes) in a natural way is also modeled on forcing semantics (sheaf semantics) (Johnstone, 1977). The variety of ways the situation of forcing appears in many different levels of mathematical reasoning (proof theory, logic, topos semantic, model theory) causes it to seem to be really fundamental phenomenon.

Forcing, despite its formal shape, has very specific action over universes of Set Theory ($\text{ZF}(\text{C})$): it defines the relations between names of things, which are not present in the Universe (it means they are not sets in this sense) such that the relations hold between sets (whose names were considered) in the enlarged Universe. The procedure of enlargement with the preservation of theorems of $\text{ZF}(\text{C})$, is just the forcing procedure. From the other perspective forcing is the recipe for proving theorems valid in enlarged Universe (which is a Model of $\text{ZF}(\text{C})$) and not necessarily valid in the ground Universe (which is also a Model of $\text{ZF}(\text{C})$). In that way Cohen was able to prove independence of Continuum Hypothesis (CH) and the Axiom of Choice (AC) from the axioms of ZF.

Bell (1983) considered the situation of forcing in some approach to Quantum Logic. He was able to point out the place in Quantum Logic, where forcing, understood logically, fails to work. There are some indications coming from his analysis that:

- (i) Set theoretical context is probably changing.
- (ii) There are some "preformal" situations where nonstandard model of reals (in the Robinsonian sense) i.e. $^*\mathbf{R}$ and standard one – $\mathbb{R}$, are "identified" (or, in fact, $\mathbb{N}$ and $^*\mathbf{N}$ as models of Peano Arithmetic (PA) are identified).

In what follows we are to investigate both points. Strikingly enough, this will lead us to rebirth of forcing in the context of QM and, in fact, forcing appears to be very intrinsic shape of Quantum Mechanical paradigm; instead of being only formal construction in logic, or QL, it is to be considered as a tool allowing the change of Metatheoretical Universe and/or the way reals appear as possible results of experiments. In fact, both points (i) and (ii) intertwine in the specific way,

which is well modeled locally by the model-theoretical constructions of Bounded Boolean Ultrasheaf of Superstructure modulo some Ultrafilter ($bBUSuS/U$) in Boolean Algebra (BA) (invented by Ozawa (1994))(see Theorem3).

In this paper we show surprising correlation of formalism of QM with some structures which are modelled on forcing constructions in various of its facets. This is still more surprising, because it was shown by Bell (1985) that the forcing formally does not work any more in QL (what we have mentioned already). The missing link for appearing forcing in QM formalism could be non classical *Main Hypo*.

3. The Rule of Indistinguishability Standard and Some Nonstandard Models of Peano Arithmetic.

The rule is exactly as stated in the title, but it requires some explanation. Firstly, the Rule cannot be formalized; otherwise it would exist formal context where one could express: Standard and Nonstandard models of PA are different and (according to the Rule) one cannot express their difference—this is impossible. For, there does exist suitably higher order formal language (in fact second order is enough), which expresses the difference between Standard $\mathbb{N}$ and Nonstandard $\mathbb{N}$; there would also exist an order, and the language of that order, which would express identity of both models. It is enough to take the language of order being maximal of both orders (with appropriate symbols), to obtain contradiction.

Secondly, the Rule is the heuristic statement only; one cannot prove it (this would require some formal context).

In what follows, we give some possible formulations of the Rule.

In some formal constructions (not necessarily connected with Nonstandard Naturals) the appearance of Nonstandard Naturals is unavoidable and out of control by formal means of the construction.

Along with the shift of the language L from the higher order into first order, we cannot distinguish Standard $\mathbb{N}$ from Nonstandard $\mathbb{N}$. There are such shifts which are not able to be described simultaneously in any formal language, which would be an extension of the language L .

*For any given formal language (of any order) there exists a pair $(\mathbb{N}, *\mathbb{N})$ such that there doesn't exist any formula $\phi(\vec{x})$ (with free variables from $\vec{x}$) in this language, which would be expressing any difference between $\mathbb{N}$ and $*\mathbb{N}$.*

An analogy with Manifolds can be usefull here: we have patches which are Standard $\mathbb{N}$ or Nonstandard $\mathbb{N}$, but the transition functions

between them cannot be formalized simultaneously with pathes (the language–logic is also to be shifted). Such a thing we call *Levelfold*.

As we will see, such a strange Rule, called from now on *Main Hypo*, has much to do with formalism of QM.

*There does not exist any formal language which would be able both to describe language of PA (counting formulas of it) and to describe **all** models of PA (There are some models of PA which cannot be described simultaneously with the formal language of PA).*

In general one can say that non simultaneous (in principle) description of formal language and the models of the theory in this language (which in fact is a core of *Main Hypo*) is analogous to non simultaneous (also in principle) measurability of some physical observables. We will see what exactly is the formal content of this coincidence.

4. Forcing Constructions.

We are ready to formulate some strange consequences of the Main Hypo. There is canonical bijection between morphisms of ultrafilters on ω and elementary embeddings of ultrapowers of $\mathbb{N}$. Morphisms of ultrafilters induce the order, which is just Rudin-Keisler (RK) order. E-morphism (End extension morphism) is induced by being an initial segment of one ultrapower of $\mathbb{N}$ in another. E-morphism also induce E-order on ultrafilters on ω . This is stronger preorder than RK. (Another possible order is the order connected to conservative extensions of ultraproducts; it is stronger than E-order. The strongest one is Rudin-Frolik order depicted as $\sqsubseteq$ (Blass, 1977). Murakami (1999) was able to define RK order and RF order on ultrafilters in arbitrarily complete BA). What follows from *Rudin – Keisler* order of ultrafilters over $P(\omega)$, taking not *Rudin – Keisler* related ultrafilters we are led to **not end extensions** of Nonstandard Naturals. Such an extension is to modify some initial segment of naturals (Nonstandard): $[0, 1, 2, \dots, \bar{n}] \subset {}^*\mathbb{N}$ in the sense that there is no 1 to 1 mapping ϕ :

$$\phi : [0, 1, \dots, \bar{n}] \xrightarrow[\text{into}]{1:1} [0', 1', 2', \dots, \bar{n}']$$

such that $\phi(n) = n'$. They both are end extensions of Standard Naturals: they have order type $\omega + (\omega + \omega^*) * \eta$, where η is dense order without endpoints and ω^* is reverse order of ω (Kamo, 1981).

Lemma 1. *Under hypothesis of non distinguishing of Standard N and some Nonstandard N (Main Hypo) it would be possible to have not end extensions of Standard Natural Numbers*

Remark: Because all extensions of PA are end extensions of initial segments of Standard Naturals (Blass, 1977), (although they could be not conservative extensions) it is clear that the above construction cannot be fully classical (see discussion in Section 3).

Lemma 2. *Under Main Hypo, one can have not end extension of Standard Naturals where some $k \in \mathbb{N}$ is modified into some infinite non standard natural from ${}^*\mathbb{N} - \mathbb{N}$.*

Next, consider lattice of projections $\mathcal{L}$ (self adjoint and idempotent operators) over some separable Hilbert space $\mathcal{H}$. Let (B_α) be some maximal BA of projections from $\mathcal{L}$.

Lemma 3. *(B_α) is complete atomic Boolean Algebra.*

Corollary 1. *Maximal algebra of projections in the lattice of projections over separable Hilbert space cannot serve as a Forcing algebra for any nontrivial forcing over any Ground Model of ZF(C).*

This is well known fact that forcing is not trivial (properly extends the ground model) iff the BA which corresponds to it is atomless BA (Jech, 1978).

Let us notice that atomic, complete BAs fulfill infinite distributive law (DL):

$$\prod_{i \in I} \sum_{j \in J} p(i, j) = \sum_{f \in J^I} \prod_{i \in I} p(i, f(i))$$

where I, J are arbitrarily, possibly infinite, sets of indices and $\{p(i, j)\}_{i \in I, j \in J}$ is any, double indexed family of elements of BA.

Any non distributive elements A, B, C from the lattice $\mathcal{L}$ (in the sense, that their spectral families are in $\mathcal{L}$):

$$(**) A \wedge (B \vee C) \neq A \wedge B \vee A \wedge C$$

cannot be simultaneously in any (B_α) . For, if not, let's consider spectral decompositions of A, B, C , from $(**)$ it follows that some triples of projections from the decompositions would also fulfill $(**)$. That means they cannot be simultaneously in any (B_α) .

Theorem 1. *(1) Under Main Hypo, there is natural correspondence between lattice of projections $\mathcal{L}$ and some family of not (ω, ω) distributive complete Boolean Algebras.*

(2) Any such Boolean Algebra can be Cohen forcing BA which adds some Cohen generic reals into some Ground model $\mathbf{V}$ of ZFC.

(3) Under Main Hypo this real interpreted in Boolean Valued model of ZFC - $\mathbf{V}^{\bar{B}}$, corresponds to the self adjoint operator which is not in (B_α) .

Theorem 2. *Cohen generic reals from Theorem1 are able to be interpreted as a probability distributions over spectrum of Boolean Algebra of Projections (B_α) in the separable Hilbert space $\mathcal{H}$. The reals might be obtained in the experiments of measuring an observable, corresponding to the self adjoint operator from Theorem1.3.*

Remark: The peculiarity of Theorem2 is in that even in the single measurement having got single "real" as a result, we already have corresponding probability distribution coming from the results of repeating measurements (to be performed). The dynamics of shifts of ZFC Models is giving this situation, which in turn enables one to "realize" *Main Hypo.*

Now we are ready to analyze what meaning can be given to reals obtained in measuring A or B (self adjoint operators). The related question is: what is the spectrum of C_ω , having fixed the spectrum of (B_α) . Both should be composed of real numbers, but once we are fixing the meaning of reals corresponding to (B_α) , then what are the reals corresponding to C_ω (interpreted, by the virtue of *Main Hypo*, in terms of projections from $\mathcal{L}$)? Finally, what does it mean that one cannot measure non commuting A and B simultaneously?.

Let us notice that reals $R^{(B_\alpha)}$ in $\mathbf{V}^{(B_\alpha)}$ and $R^{((B_\alpha)+(A_s))} = R^{C_\omega}$ in $\mathbf{V}^{C_\omega}$ differ due Cohen generic reals on the level of ground model $\mathbf{V}$. This means that we have to change the ground model along with the shift $(B_\alpha) \rightarrow ((B_\alpha) + (A_s))$, where both algebras are treated as algebras of projections from $\mathcal{L}$. That means in turn, that to maintain indistinguishability of algebras, we have changed the meaning of the Lattice of Projections. Thus we have reached the point where everything has changed the meaning (even standard Natural Numbers from *Main Hypo*).

To try to see the difference between varying Naturals or Reals we trying to fix $\mathbb{N}$ and $\mathbb{R}$ and are going to employ the technique of Bounded Boolean Ultrasheaf of Superstructure Modulo some Ultrafilter in BA (*bBUSuS/U*) (Ozawa, 1994). *bBUSuS/U* is in fact based on generalization of Ultraproduct modulo Ultrafilter in $P(\omega)$ (Keisler et al., 1990) into Boolean Ultraproduct modulo Ultrafilter in BA in question (Mansfield, 1971). The next step is to construct Superstructure to generate nonstandard elements inside Boolean Valued Universum of ZFC, and to make forcing, and to generate infinitesimals simultaneously by the same Ultrafilter in BA (Ozawa, 1994).

On the base of fixed $\mathbb{R}$ we will be able to see what reals should look like in the presence of non compatible observables. Let us consider some superstructure allowing one to deal with $\mathbb{N}$, $^*\mathbb{N}$, $\mathbb{R}$ and $^*\mathbb{R}$ according to some Ultrafilter on $P(\omega)$ i.e. $V(\mathbf{R} \cup P(\omega))$. On the side of "true"

(connected to Standard $\mathbf{N}$ being well distinguished to any Nonstandard $\mathbf{N}$) Lattice of Projections, maximal BA of projections in $\mathcal{L}$ is isomorphic to $P(\omega)$, (Lemma3). Following the arguments in Ozawa (1994) we can define superstructure in Boolean valued model $\mathbf{V}^{P(\omega)} \simeq \mathbf{V}^{(B_\alpha)}$ (being superstructure is expressible by Δ_0 - formulas). This means that our superstructure generates $^*\mathbf{N}$ and $^*\mathbf{R}$ which are equally generated inside $\mathbf{V}^{(B_\alpha)}$.

By the shift $(B_\alpha) \rightarrow ((B_\alpha) + (A_s)) \simeq C_\omega$ (coming from *Main Hypo*), we are faced with Boolean valued model $\mathbf{V}^{C_\omega}$ and ingeneric reals are appearing with respect to the superstructure $V(R \cup C_\omega)$ (Ozawa, 1994). Now, there is nonequivalence between $^*\mathbf{N}$, $^*\mathbf{R}$ generated in the outside and inside of the $\mathbf{V}^{C_\omega}$. More exactly, an effect of forcing corresponds inside $\mathbf{V}^{C_\omega}$ to ingeneric reals with respect to fixed $\mathbf{R}$. That means, *Main Hypo* through the shift $\mathbf{V}^{P(\omega)} \simeq \mathbf{V}^{(B_\alpha)} \rightarrow \mathbf{V}^{C_\omega}$ is giving ingeneric reals as those which correspond to enlargement $(B_\alpha) \rightarrow ((B_\alpha) + (A_s))$, so ingeneric reals correspond to A (A determines generic Cohen real, which correspond to some ultrafilter U in C_ω . C_ω and U in turn, give ingeneric reals (Ozawa, 1994)). Let us remind that by a spectrum of Boolean Algebra of projections we mean the spectrum of C^* - algebra, which is uniquely determined by the algebra of projections (Jech, 1984). Now we see that fixing the meaning of true $\mathbf{R}$ as connected to the spectrum of (B_α) we can regard reals connected to the shifted algebra $((B_\alpha) + (A_s))$ as ingeneric enlargement coming from $bBUSuS/U$ with respect to C_ω and ultrafilters in it. We have obtained the characterization of the spectrum of enlarged algebra coming from non compatible observables (through *Main Hypo*).

Theorem 3. *In the measurement of non compatible observables as a result of the possible simultaneous measurement we would have real number and some ingeneric real corresponding to Bounded Boolean Ultra-sheaf of SuperStructure modulo some Ultrafilter in BA — $bBUSuS/U$. That is why having both reals is not possible.*

Theorem 4. *According to the shift $(B_\alpha) \rightarrow ((B_\alpha) + A)$ we have the shift, corresponding to the forcing in the ground model: $\mathbf{V} \rightarrow \mathbf{V}[r_A]$, where r_A is Cohen generic real corresponding to A .*

Proof: $\mathbf{V} \simeq \mathbf{V}^{(B_\alpha)}/U \rightarrow \mathbf{V}^{((B_\alpha)+A)}/\bar{U} \simeq \mathbf{V}^{C_\omega}/\bar{U} \simeq \mathbf{V}[r_a]$. $\square$

Remark: In Theorem4 we cannot have both real numbers as a result of measuring non compatible observables, otherwise they would be compatible, and an enlarged the algebra $((B_\alpha) + (A_s)) \simeq C_\omega$ does not appear. We only have algebra (B_α) and consequently standard reals as a result of measuring observables from this algebra.

In that way we have obtained the model of being non compatible observables via forcing construction, in the sense that we are able to

show what spectrum could correspond to the "algebra" generated by non compatible observables. Let us note that our forcing involved in the construction is one which changes the meaning of the Boolean Valued universe as in $bBUSuS/U$.

Corollary 2. *Under Main Hypo the spectrum of the Boolean algebra (B_α) is shifted to the spectrum of enlarged Boolean algebra $((B_\alpha)+(A_s))$ by adjoining some ingeneric reals.*

Now we are interested in the characterization of the meaning of standardness of real numbers (and standardness of related set-theoretical entities) appearing in measurement of various observables; what are our formal abilities to decide weather we can maintain the notion of standard real over all class of observables? The above construction allowed us to "see" Nonstandard reals or Ingeneric reals just by referring to well defined notion of Standard Natural or Standard Real. Is it possible to maintain it for all observables and measurements?

It can be shown that this is not the case; the modification of Standard Naturals and reals has to be taken seriously in the context of QM. This means we cannot decide once and for ever about the standardness of sets to "see" (with the bounded Boolean ultrasheaf constructions) the shifts Standard Reals $\rightarrow$ Ingeneric Reals; this is possible only locally.

To close this chapter, let us make a note about some possibilities to consider "Spectral Theorem" associated to the "Boolean algebra", that come from non compatible observables, as in Theorem2. Firstly, there is well developed theory of measure on the internal nonstandard spaces (Loeb, 1982), so called Loeb measure. Secondly, Ozawa (1994) has extended this measure over ingeneric measure spaces. We can try to characterize " C^* -algebra" coming from the above "Boolean algebra" generated by non commuting observables, as the one, which in some sense is isomorphic to the "function algebra" over ingeneric spectrum. In this way, noncommutativity is eaten by our *Main Hypo* and the commutative function algebra emerges again.

5. Model Theoretic Analysis of Small Exotic Smooth Structures on $\mathbb{R}^4$.

General references on exotic smoothness topics and Kirby calculus are Gompf et al. (1999) or Kirby (1989). There were also attempts to relate exotic smoothness to some physical valid situations (Brans, 1994, Śladowski, 2001). Let us consider so called small exotic smooth R^4 . Small exotic smooth R^4 's are all imbedded into Standard $\mathbb{R}^4$. They come from the failure of smooth h-cobordism theorem in dimension 4

(large exotic smooth structures on $\mathbb{R}^4$ come from the failure of higher dimensional surgery theory applied to dimension 4). There are continuum many different non diffeomorphic smooth structures (small as well large) on topological $\mathbb{R}^4$. Any large smooth R^4 contains compact manifold which can not be smoothly imbedded into standard $\mathbb{R}^4$. It is known that small exotic smooth R^4 , as being 4-manifolds, have handle decompositions (infinite) with only 0, 1 and 2-handles included. Typical handlebody of such manifold is given by a compact submanifold of $\mathbb{R}^4$ (so called Akbulut cork) with some Casson handles (CH) attached. CH are infinite towers of kinky handles and any Casson handle is homeomorphic to the standard 2-handle (Freedman, 1982). The simplest exotic (small) R^4 is represented by the following Kirby diagram (Biřaca et al., 1996)(Figure 1), where it is understood that we are taking only interior of the handlebody ignoring the boundary (except the attaching circle of the CH). Different exotic small R^4 differ among themselves just by the complexity of CH and by compact K .




Figure 1. Casson handle in the simplest exotic R^4 .

The motivation to consider small exotic R^4 is in that, they might be represented (in principle) by the explicit figures with their Kirby diagrams. The idea to use Model Theory to analyze exotic structures on $\mathbb{R}^4$ is in our ability to introduce specific partial order in the set of pairs $(K, CH_i^{(n)})$ of compacts K and finite stages of Casson Handles (attached to K to obtain exotic smooth R^4). The partial order in question is to be separative one. Then we can produce boolean algebra of regular open (RO) subsets of the above partial order P . So, we have $RO(P)$ (Jech, 1978). It is known that this algebra is complete atomless BA (Jech, 1978). Taking Ground Model V (as usual this is to be Countable Transitive Model of ZFC to allow forcing), we can create Boolean valued model of ZFC – $V^{RO(P)}$. Based on this, we are able to retrieve two valued models just by taking quotients by non principal

ultrafilters in $RO(P)$. On the level of Ground Model this is just the same as adding some Cohen generic real $r: V \rightarrow V[r]$.

Now we can formulate:

Theorem 5. *Forcing which adds generic Cohen real r to V corresponds exactly to attaching some Casson Handle to K , corresponding in turn to the change of the smooth differential structure on $\mathbb{R}^4$.*

Next, we are ready to describe how the *Main Hypo* is going to generate Smooth Exotic Structures on $\mathbb{R}^4$.

Lemma 4. *Under Main Hypo one can have indistinguishability of some Standard $\mathbb{R}^4$ and some Non Standard ${}^*\mathbf{R}^4: \mathbb{R}^4 \approx {}^*\mathbf{R}^4$.*

Lemma 5. *Dirac δ -function (distribution) on $\mathbb{R}^4$, is smooth ordinary function (with values in ${}^*\mathbf{R}$) on ${}^*\mathbf{R}^4$.*

Proof: see Robinson (1979).

Now, let us observe, that having $\mathbb{R}^4 \approx {}^*\mathbf{R}^4$ we do not distinguish formally between smooth functions on ${}^*\mathbf{R}^4$ and smooth functions on $\mathbb{R}^4$. This means we should have some smooth function on $\mathbb{R}^4$ which corresponds (not in a direct way) to δ -function on $\mathbb{R}^4$. The change in the smoothness which guarantees the change δ -function $\rightarrow$ smooth function, is, of course, the same as the changing standard smoothness into some other one (nondiffeomorphic). This is Non Standard Exotic Smooth Structure.

We have reached the point where Model Theoretical analysis of exotic smooth structures on $\mathbb{R}^4$ shows connections to Set Theoretical forcing and *Main Hypo*. Is it accidental, or does it express any essential correlations between Exotic Smooth $\mathbb{R}^4$ and Model Theoretical shape of QM (which we have exhibited in the first part)? We claim that the former is true, but detailed analysis will appear elsewhere.

In what follows we only give an example concerning this situation in the context of the so called *AdS/CFT* correspondence in *Sustring* or M-theory.

6. Model Theoretic Analysis of the *AdS/CFT* correspondence – an example.

Since Maldacena (1997) made the conjecture about strict mathematical equivalence (duality) of completely different theories: Superstring Theory on $AdS_5 \times S^5$ background and Supersymmetric Conformal $SU(N)$ Yang Mills Theory in 4-dimensions in the limit of large N , much work has been done to check it. There were also some proposals

(Witten et al., 1999) to make connection of this duality to "realistic" non supersymmetric, non conformal YM theory in 4-dim (QCD?).

Based on some Model Theoretic analysis of the solutions of 10-dim Supergravity equations we present possible mechanism to generate the sources of masses terms in 4-dim Super YM CFT which might (in principle) break SuSy.

The crucial ingredient of AdS/CFT correspondence is to take N D3-branes as a solutions of classical SuGry equations in 10-dimensions (which is low energy limit of IIB SuString Theory and D3-branes are also present here). Taking N to infinity and considering N D3-branes as coincident, it is possible to recover the near horizon geometry of stack of D3-branes so obtained. This geometry is just $AdS_5 \times S^5$. Witten (1998) has shown how the correlation functions in the bulk SuString Theory correspond to the ones in SYM CFT on the 4-dim boundary of AdS_5 . He has also shown what is the shape of the sources in boundary theory (coming from the bulk) and to what boundary operators they couple. We don't present calculations here, we just state main ideas about how one might generate sources of gravity which would break SuSy in the bulk YM theory.

Let us consider the stack of N D3-branes as a multiplicity of N 4-dim world volumes which is (topologically) a multiplicity of $\mathbb{R}^4$. Let us note that every D3-brane is the solution of 10-dim SuGry. This can be seen as a kind of model of the theory given by 10-dim SuGry equations. We are going to produce another model (Nonstandard) using technique of Ultraproduct (modulo some ultrafilter in $P(\omega)$). We don't discuss here the subtleties concerning orders of the formal languages involved. In such a way we will generate Nonstandard solution to 10-dim SuGry whose world volume is just $(\mathbb{R}^4)^{N \rightarrow \infty} / U$ which is ${}^*\mathbf{R}^4$. Now we are able to state:

Theorem 6. *Under Main Hypo the procedure described above gives sources of gravity in 4-dim.*

The proof of this Theorem is based on the analysis in the previous paragraph, where the shift δ -function $\rightarrow$ smooth function on $\mathbb{R}^4$, corresponded to the change of differential structure on $\mathbb{R}^4$, and we know from the work of Asselmeyer (1997), how the change of smooth differential structure on compact 4-Manifold generates sources of 4-dim Gravity. The possibility of breaking SuSy by these sources comes from analysis of Polchinski and Strassler (Polchinski et al., 2000). The details are to be presented elsewhere.

7. Summary

We have presented, very in a sketchy way, the ideology of analyzing QM via Model Theory based on *Main Hypo*. Even more sketchily we have noticed about similar analysis of the exotic (small) smooth structures on $\mathbb{R}^4$. There are striking similarities to both approaches: they require *Main Hypo* and a set theoretical forcing appears unexpectedly, yet rather naturally. *Main Hypo* and forcing are able to code formally the structure of QM in the language of operator algebras over separable Hilbert space. This coding can be used to explain certain aspect of Maldacena conjecture that quantum field theory without gravity is to be dual to the theory describing gravity.

There are also some possibilities to break SuSy or conformal invariance in 4-dim YM Theory in order to obtain connections to realistic (confined) 4-dim QCD.

References

- Albeverio S., Fenstad J. E., R. Høegh-Krohn Lindstrøm T. (1986). *Nonstandard Methods in Stochastic analysis and mathematical Physics*, Academic Press, Orlando.
- Asselmeyer T. (1997). Generation of Source Terms in General Relativity by Differential Structures, *Classical and Quantum Gravity* **14**, 749-758.
- Bell J. L. (1983). *Orthologic, Forcing, and the Manifestation of Attributes* Suthest Asian Conference on Logic, eds. C. T. Chong and M. J. Wicks, Elsevier Science Publishers, North-Holland.
- Bell J. L. (1985). *Boolean-Valued Models and Independence Proofs in Set theory*, 2nd ed. Oxford University Press, Oxford.
- Bell J. L. (1985). Orthospaces and Quantum Logic, *Foundations of Physics* **15**, no 12, 1179-1202.
- Benioff P. (2001). quant-ph/0103078, quant-ph/0106153.
- Bižaca Ž., Gompf R. (1996). Elliptic Surfaces and Some Simple Exotic R^4 's, *Journal of Differential Geometry* **43**, 458-504.
- Blass A. (1977). End Extensions, Conservative Extensions, and the Rudin-Frolik Ordering, *Transactions of the American Mathematical Society* **225**, 325-340.
- Brans C. H. (1994). *Journal of Mathematical Physics* **35**, 5494.
- Cohen P. J. (1966). *Set Theory and the Continuum Hypothesis*, Benjamin, New York, 1966.
- Dalla Chiara M., Giuntini R. (2001). *Quantum Logics*, in Handbook of Philosophical Logic, ed. D. Gabbay, Kluwer Dordrecht.
- Fearins J. D. (2002). *A Physical Quantum model in a Smooth Topos*, quant-ph/0202079.
- Freedman M. H. (1982). The Topology of Four-Dimensional Manifolds, *Journal of Differential Geometry* **17**, 357-453.
- Gompf R. E., Stipsicz A. I. (1999). *An Introduction to 4-Manifolds and Kirby Calculus*, American Mathematical Society, Rhode Island.
- Guts A., Zvyagitser A. (1999, 2001). physics/9909016, gr-qc/0001076.

- Isham C., Butterfield J. (1999). *International Journal of Theoretical Physics*, **38**, 827.
- Jech T. (1978) *Set Theory*, Academic Press.
- Jech T. (1984). Abstract Theory of Abelian Operator Algebras: An Application of Forcing, *Transactions of the American Mathematical Society* **289**, no 1, 133-162.
- Johnstone P. T. (1977). *Topos Theory*, Academic Press Inc., London.
- Kamo V. (1981). Nonstandard Natural Number System and Nonstandard Models, *Journal of Symbolic Logic* **46**, 365-376.
- Keisler H. J., Chang C. C. (1990). *Model Theory*, 3rd ed., North-Holland, Amsterdam.
- Kirby R. C. (1989). The Topology of 4-manifolds, *Lectures Notes in Mathematics* **1374**, Springer-Verlag, Berlin, Heidelberg, New York.
- Kochen S., Specker V. (1967). *Journal of Mathematics and Mechanics* **17** No 1.
- Kock A. (1981). *Synthetic Differential Geometry*, London Mathematical Society Lecture Notes, Series **51**, Cambridge U. Press.
- Król J. (2001). Formal Languages and Model Theoretic Perspectives in Physics, *Acta Physica Polonica B* **32** No 11, 3855-3871.
- Król J. (2002) Problem of Background Independence in Quantum Gravity and Forcing Constructions, *in preparation*.
- Loeb P. A. (1982). *Measure Spaces in Nonstandard Models underlying Standard Stochastic Processes* Proceedings of the International Congress of Mathematics held in Warsaw, PWN, North Holland **1**, 323-336.
- MacLane S., Moerdijk I. (1992). *Sheaves in geometry and Logic: A First Introduction to Topos Theory*, Springer-Verlag, 1992.
- Maldacena J. (1997). *Adv. Theor. Math. Phys.* **2**, 231, [hep-th/9711200].
- Mansfield R. (1971) The Theory of Boolean Ultrapowers, *Annals of Mathematical Logic* **2**, 297-323.
- Reyes G. E., Moerdijk I. (1991). *Models for Smooth Infinitesimal Analysis*, Springer-Verlag, New York.
- Murakami M. (1999). Standardization Principle of Nonstandard Universes, *Journal of the Symbolic Logic* **64**, No 4, 1645-1655.
- von Neuman J., Birghoff G. (1936). *Annals of Mathematics* **37**, 823.
- Ozawa M. (1994). Forcing in Nonstandard Analysis, *Annals of Pure and Applied Logic* **68**, 263-297.
- Polchinski J., Strassler M. J. (2000). The String Dual of a Confining Four-Dimensional Gauge Theory, hep-th/0003136.
- Raptis I. (2001). Preasheaves, Sheaves and their Topoi in Quantum Gravity and Quantum Logic, gr-qc/0110064.
- Robinson A., Barwise J. (1971). Completing Theories by Forcing, *Annals of Mathematical Logic*, **2**, 119-142.
- Robinson A. (1966). *Non-Standard Analysis*, Studies in Logic and the Foundations of Mathematics, Amsterdam.
- Robinson A. *Topics in Non-Archimedean Mathematics*, Collected Papers, 99-112.
- Scott D. (1969). *Boolean Models and Nonstandard Analysis*, in Applications of Model Theory to Algebra, Analysis, and Probability, Holt, Reinhart, Winston, New York, ed. W.A. J. Luxemburg.
- Rasiowa H., Sikorski R. (1963). *The Mathematics of Metamathematics*, Warszawa.
- Śladowski J. (2001). Gravity on Exotic R^4 's with few symmetries, *International Journal of Modern Physics D* **10**, 311.
- Takeuti G. (1978). *Two Applications of Logic to Mathematics*, Mathematical Society of Japan **13**, Kano Memorial Lectures 3.

- Takeuti G. (1979). *Boolean Valued Analysis* Applications of Sheaves, eds. Fourman, Mulvey, Scott, Lectures Notes in Mathematics **753**, Springer, Heidelberg, 714.
- Takeuti G. (1983). *Quantum Set Theory* in Current Issues in Quantum Logic, eds. E. Beltrametti and B. C. van Frassen, Plenum New York, 303.
- Vopenka P. (1965). The Limits of Sheaves and Applications on Constructions of Models, *Bulletin de l'Academie Polonaise des Sciences* **13**, 189-192.
- Witten E., Klebanov I. R. (1999). AdS/CFT Correspondence and Symmetry Breaking, *Nuclear Physics B* **556**, 89-114.
- Witten E. (1998). Anti-de Sitter space and holography, *Advances in Theoretical and Mathematical Physics* **2**, 231, [hep-th/9802150].